

***Harpephyllum caffrum* stimulates glucose uptake, Abates Redox Imbalance and, Modulates Purinergic and Glucogenic Enzyme Activities in Oxidative Hepatic Injury**

Kolawole A. Olofinisan<sup>1</sup>, Ochuko L. Erukainure<sup>2</sup>, Beseni K. Brian<sup>1</sup>, Md. Shahidul Islam<sup>1</sup>

1. Department of Biochemistry, School of Life Sciences, University of KwaZulu – Natal, (Westville Campus), Durban 4000, South Africa
2. Department of Pharmacology, University of the Free State, Bloemfontein 9300, South Africa.

**Corresponding author:** [islamd@ukzn.ac.za](mailto:islamd@ukzn.ac.za); Telephone: +27312607942

**Supplementary Table 1.** IC<sub>50</sub> values of *Harpephyllum caffrum* infusion in *in vitro* biological assays (µg/mL).

	Infusion	Trolox	Acarbose	Orlistat
FRAP	47.9	74.1	ND	ND
OH <sup>•</sup> scavenging	32.3	> 320	ND	ND
DPPH scavenging	45.2	> 320	ND	ND
α-amylase	298.2	ND	> 320	ND
α-glucosidase	5.6	ND	206.7	ND
Lipase	88.8	ND	ND	40.9

ND = Not determined.