

Supplementary Data

Dieckol isolated from *Eisenia bicyclis* extract suppresses RANKL-induced osteoclastogenesis in murine RAW 264.7 cells

Running title: Osteoclastogenesis inhibitory activity of dieckol

Su-Hyeon Cho^{1,2†}, Tae-Hyung Kwon^{3†}, Hoibin Jeong¹, Jin Sook Kim¹, Song-Rae Kim¹, Myeong Seon Jeong¹, SeonJu Park¹, Miri Choi¹, Jung-Hee Woo⁴, Juhee Ahn^{2*}, Kil-Nam Kim^{1,5*}

¹ Chuncheon Center, Korea Basic Science Institute, Chuncheon 24341, Republic of Korea

² Department of Medical Biomaterials Engineering, College of Biomedical Sciences, Kangwon National University, Chuncheon 24341, Republic of Korea

³ Department of Research and Development, Chuncheon Bio-industry Foundation, Chuncheon, Republic of Korea


⁴ Marine Industry Research Institute for East se rim, Uljin-gun, Gyeongsangbuk-do, Republic of Korea

⁵ Department of Bio-analysis Science, University of Science & Technology, Daejeon, 34113, Republic of Korea


†: Both authors contributed equally to this work.

* Corresponding author: E-mail: knkim@kbsi.re.kr (K.N. Kim); juheeahn@kangwon.ac.kr (J. Ahn)


(A)




(B)



(C)



(D)



Supplementary Figure 1. Chemical structures of (A) 6,6'-bieckol, (B) 6,8'-bieckol, (C) 8,8'-bieckol, and (D) dieckol.