

# Anthrahydroquinone-2,6-disulfonate alleviates paraquat-induced kidney injury *via* the apelin-APJ pathway in rats

Qi Li<sup>1#</sup>, Bo Wang<sup>2#</sup>, Kai-Wen Lin<sup>3</sup>, Tang Deng<sup>1</sup>, Qi-Feng Huang<sup>1</sup>, Shuang-Qin Xu<sup>1</sup>, Hang-Fei Wang<sup>1</sup>, Xin- Xin Wu<sup>1</sup>, Nan Li<sup>1</sup>, Yang Yi<sup>1</sup>, Ji-Chao Peng<sup>1</sup>, Yue Huang<sup>1</sup>, Jin Qian<sup>1✉</sup>,  
Xiao-Ran Liu<sup>1✉</sup>

<sup>1</sup>*College of Emergency and Trauma, Hainan Medical University, Key Laboratory of Emergency and Trauma of*

*Ministry of Education, The First Affiliated Hospital of Hainan Medical University, Haikou 571199, China*

<sup>2</sup>*Jiangdong Health Center, Meilan District, Haikou 571126, China*

<sup>3</sup>*Hainan Women and Children's Medical Center, Haikou 570312, China*

✉To whom correspondence may be addressed. E-mail: 798052058@qq.com (J. Qian);

hy0203049@hainmc.edu.cn (X.R Liu).

**Table 1. Renal histopathological scoring standard.**

The mean score of 12 visual field	Injury degree	Ratio of abnormal visual field
0	Normal	Normal nephron and stroma
1	Light	< 25%
2	Moderate	25-50%
3	Severe	> 50%